



Cooling capacity of urban trees exposed to thermal stress



Luiza Maria Teophilo Aparecido¹

Kallika Naylor^{2*}, Isabella Werner³, Raven Braithwaite¹, Ella Schmidt⁴, Heather Throop^{1,2}, Jnaneshwar Das¹, Kevin R Hultine⁵

¹School of Earth and Space Exploration, Arizona State University, Tempe, AZ, USA – lmtephilo@gmail.com

²School of Life Sciences, Arizona State University, Tempe, AZ; ³Ira A. Fulton Schools of Engineering, Arizona State University, Tempe, AZ; ⁴Paradise Valley High School, Phoenix, AZ; ⁵Desert Botanical Garden, Research, Conservation, and Collections, Phoenix, AZ

Background

Urban heat islands pose a major threat to human and environmental health, requiring city managers to develop strategies to mitigate the effects of excess heat. Urban vegetation is known for mitigating some of these threats by cooling the local surroundings through shade and evapotranspiration. However, choosing the appropriate tree species remains a challenge, as urban planners have little information on species' cooling capacities (i.e., shade intensity and area, water use).

Objectives

We quantified the species-specific cooling effect of 14 tree species (native and exotic) commonly found in Phoenix, AZ, USA. Specifically:

- 1) How does plant water loss (stomatal conductance, g_{sw}) and transpiration, (Et) vary across urban tree species (native vs. exotic)?
- 2) How did these parameters change from spring to summer, when temperatures regularly exceed 40 °C?

Methods

Research Site: Desert Botanical Garden, Phoenix, AZ, USA

Temporal extent: April-July, 2021. Measurements taken 4-5 days at the end of each month

Sample plants: 14 species chosen based on Maricopa County Urban Tree Selection Criteria list.

- 15-gal potted plants from local nurseries irrigated daily (once in spring, twice in summer). Four plants per species

Measurements:

- **Diurnal gas exchange:** porometer (LI600), and photosynthesis system (LI6400) for small leaves and needles → 4 leaves per plant
- **Thermal:** thermal IR sensor, FLIR One phone camera
- **Weather:** nearby HOBO weather station
- **Plant traits:** LMA, LDMC, thermal damage, canopy area and volume




1 Description of sampled plants and percentage difference in stomatal conductance (Δg_{sw}) and transpiration (ΔEt) from spring (April-May) to summer (June-July).


Code	Species	Common Name	Origin	Δg_{sw}	ΔEt
Babl	<i>Bauhinia blakeana</i>	Hong Kong Orchid tree	Exotic	24%	52%
Chli	<i>Chilopsis linearis</i>	Desert willow	Native	75%	84%
Cuar	<i>Cupressus arizonica</i>	Arizona cypress	Native	42%	49%
Frve	<i>Fraxinus velutina</i>	Arizona ash	Native	24%	46%
Juma	<i>Juglans major</i>	Arizona walnut	Native	-8%	-1%
PaDM	<i>Parkinsonia x 'Desert Museum'</i>	Palo verde	Native	31%	26%
Pibr	<i>Pinus brutia</i>	Eldarica pine	Exotic	61%	62%
PIRP	<i>Pistacia x 'Red Push'</i>	Chinese pistache	Exotic	9%	34%
Plwr	<i>Platanus wrightii</i>	Arizona sycamore	Native	-62%	-5%
Prgl	<i>Prosopis glandulosa</i>	Honey mesquite	Native	63%	64%
Quvi	<i>Quercus virginiana</i>	Live oak	Exotic to AZ	-26%	15%
Sela	<i>Searsia lancea</i>	African sumac	Exotic	-26%	15%
Ulpa	<i>Ulmus parvifolia</i>	Chinese elm	Exotic	-21%	19%
Vaca	<i>Vauquelinia californica</i>	California rosewood	Native	-23%	21%



2 Average transpiration (A) and stomatal conductance (B) per species. Solid black enlarged points indicate exotic species.




3 Climatic (A) and plant water use (B-C) variation across measurement months. Letters indicate significant differences ($\alpha=0.05$) of transpiration and stomatal conductance between months.



Results

- Et rates increased 33% from spring to summer; and g_{sw} increased 12%.
- Mean g_{sw} was negatively affected by both VPD_{leaf} and T_{leaf} . Mean Et kept constant (ns), with exception to negative response to VPD_{leaf} during June/2021.
- Wide range of Et rates reflects different strategies that each species adopt to cope with heat (evaporative cooling vs. stomatal suppression).
- Mean Et and g_{sw} (marginally) were both negatively correlated only to LMA. Relationship to LMA was strongest during the summer months.
- T_{diff} became mostly positive in July/2021 after the heatwave, suggesting stomatal closure from hydraulic failure or leaf thermal damage. Highest T_{diff} was in June/2021 indicating that stomatal regulation was focused on evaporatively cooling the canopy and surroundings.

4 Effects of leaf temperature on transpiration (A) and stomatal conductance (B) across species; effects of temperature difference (T_{diff}) on transpiration (similar to g_{sw}) across months (C); relationship to leaf mass per area (D).



Conclusions

- *Pinus eldarica*, *Cupressus arizonica*, *Chilopsis linearis*, *Searsia lancea*, and *Fraxinus velutina* were the most "cooling efficient" trees for Phoenix landscaping → provide intermediate-full shade across seasons, relative to water use under arid conditions.
- Highly drought-tolerant plants used the most water and partially dropped leaves during the summer (*Quercus*, *Parkinsonia*, and *Prosopis*).
- **In progress:** NDVI variation across seasons and species; leaf water use upscaled to whole-canopy.

Acknowledgements

This presentation contains partial findings from Kallika Naylor's undergraduate thesis entitled "Water use of desert urban tree species across seasons" defended on 6 Dec 2021.

- Research funding: ASU SESE Exploration Postdoctoral Fellowship awarded to Dr. Aparecido.
- Student scholarships awarded through ASU/NSF WAESO (Western Alliance to Expand Student Opportunities).
- Special thanks for the support given by the Desert Botanical Garden, Throop lab, Das lab and Hultine lab.

✪ **Next steps:** I will be starting my lab as an Assistant Professor at Univ of Utah in July/2023. Please reach out if you are interested in collaborating in future projects related to this work.