



Hemiptera: Lygaeidae
Photo by J. Patterson

Effects of habitat type on arthropod community structure in a heterogeneous urban environment

Nancy McIntyre, Jessamy Rango, William Fagan, & Stan Faeth
Arizona State University




Coleoptera: Meloidae
Photo from Colorado State University

Research objectives


- describe composition and turnover of arthropod communities in 4 types of urban land use in the Phoenix, Arizona, metropolitan area
- determine which taxa are indicative of the area's dominant forms of urban land use
- explore how variation in physical habitat structure may explain variation in arthropod communities

Land use: Phoenix metropolitan area



Arthropod collecting sites were chosen to represent the 4 most dominant forms of land use in Phoenix.

Methods: long-term arthropod monitoring



Methods: physical habitat structure


Percent ground cover of each of the following was measured in a 25-m-diameter circle, centered around each site:

- | | | |
|-------------|---------------|-------------------|
| buildings | native trees | grass |
| bare ground | native shrubs | herbaceous |
| gravel | exotic trees | agricultural crop |
| concrete | exotic shrubs | |
| rock | cactus | |


Overall results

- Arthropods from 88 taxa (21 orders, 65 families) were collected. Ants, mites, and springtails were ubiquitous and accounted for 93% of individuals captured.
- As many taxa were found in agricultural and residential lands as in native desert. The fewest taxa were found at industrial sites.
- Taxonomic richness tracked temperature more closely than it did precipitation.
- Differences in community composition among land-use types corresponded to differences in habitat structure with land use.


Results: arthropod richness



Results: community composition




Results: seasonal richness with climate variables



Results: habitat structure and community composition

Canonical Correspondence Analysis



- = abundance of each taxon (n=88 taxa)
- ➔ = linear combinations of physical habitat features: arrow length = importance of habitat feature, angle between arrows = correlation (small angle = high correlation)
- most anthropogenic habitat features are situated in lower portion of graph
- the abundance of points clustered around origin indicate that most taxa are habitat generalists
- named taxa illustrate patterns of arthropod abundance as related to habitat structure

Implications

- the presence of spatial heterogeneity within the Phoenix metro area boosts the overall arthropod diversity of the region
- although the number of taxa is similar among land-use types, the community composition differs, reflecting differences in the physical habitat structure with land use
- there are arthropod communities that are characteristic of different forms of urban land use, which may be very useful in detecting latent effects of future urban development

Acknowledgments

Jim Barnette
Dave Boomgaard
Rob Brandt
Sharlene Cardona
Joelle Don de Ville
Michelle Fink
Gerry Foster
Marc Hinze
Chris Lawrence
Vickie Massey
Lisa McKelvy
Gary Patterson
Darlene Sitzler
Maggie Tseng
Sean Walker



Coleoptera: Coccinellidae
Photo by N. McIntyre