

### A Study of Organic Compounds in Haboob Particulate Matter

ARIZONA STATE UNIVERSITY

Jershon Eagar, Denise Napolitano, Aurelie Marcotte, Anthony Anderson, Joana Sipe, Blanca Rodriguez, and Pierre Herckes

Department of Chemistry and Biochemistry, Arizona State University, Tempe, AZ


### Introduction

Each July marks the beginning of the summer monsoon season in the southwestern United States. In Phoenix, thunderstorms are frequently preceded by dust storms, or haboobs. Haboobs suspend sand and dust in the air, allowing them to commingle with urban particulate matter (PM).

The chemical composition of haboob dust is not well known and it remains unclear whether the composition of haboob dust significantly impacts background urban PM. Comparisons of the carbonaceous compositions of background urban PM and haboob PM can provide useful information regarding potential mutual sources as well as potential scavenging and agglomeration activity during dust storms.

## Weather Conditions (US NOAA) Lat: 33.427799 Lon: -112.003465 Elev: 340m Phoenix Sky Harbor International Airport 100 880 20 28-Jun 8-Jul 18-Jul 28-Jul 7-Aug 17-Aug 27-Aug 6-Sep 16-Sep 26-Sep Date — Relative Humidity (%) — Wind Speed (mph) — Gust Speed (mph) — Visibility (miles)

### **Photos**





**Above**: View of high-volume samplers the day following a rain and thunder storm.

### **Experimental Procedures**

### **Sample Collection**


Haboob PM and background urban PM samples were collected during the July 2013 – September 2013 monsoon season at the ASU campus in Tempe, Arizona using high-volume aerosol samplers.

### Analysis


PM2.5 fractions (PM with aerodynamic diameters less than 2.5 micrometers) were analyzed by thermal optical transmittance (TOT) to determine differences in atmospheric concentrations of organic and elemental carbon in background urban PM and haboob PM. The samples were then analyzed by GC/MS to identify polycyclic aromatic hydrocarbons (PAH). The PM>2.5 fractions (PM with aerodynamic diameters greater than 2.5 micrometers) were analyzed by TOT to determine differences in atmospheric concentrations of total carbon in background urban PM and haboob PM.

### **Weather Data**

Weather and air quality data were obtained from NOAA and the US EPA Air Quality System at two locations close to the Tempe ASU campus (see map below).



# Temporal Trends of Haboob PAH in PM2.5 25 20 Perylene Perylene 10 8 /80/95720M4 to uotatata and the second of the second o



### Legend ↓ Rain ↓ Haboob (Dust storm) \* Haze

### **Acknowledgments**

We would like to thank the EPA for providing the Air Quality System (AQS) Data Mart and the Technology Transfer Network (TTN).

### **Main Observations**

Haboob events are characterized by short spikes in PM2.5 and PM>2.5 concentrations. Following the event, concentrations go down to typical summer urban levels rather quickly. The PM2.5 shows small spikes during the Haboob events since most of the dust is in larger size fractions.

Elemental (EC) and Organic carbon (OC) are significant components on fine particulate matter (PM2.5) in Phoenix, AZ (approximately 10 – 60 %).

The relative contributions of OC and EC in PM2.5 decrease during most Haboob events, an indicator that most Haboob aerosol material is inorganic or non-carbonaceous, consistent with soil components.

The total carbon concentrations in PM>2.5 is low (often about 1ug/m³; data not shown). This is consistent with larger particulate matter consisting of mainly mineral soil dust. The PM>2.5 also did not change significantly during Haboob events.

Perylene, a 5 ring PAH from combustion sources and an example organic compound, is ubiquitous in urban air. Its concentration decreases during most Haboob events but then quickly goes back to typical levels. However, when hazy conditions preceded a dust event (annotated by ♣), the concentration of Perylene and PM2.5 increased significantly (74pg/m³ and 22ug/m³ respectively). Such an increase can be linked to resuspension of road and urban dust containing PAHs.

Rain events can also impact carbonaceous aerosol but did not seem to have altered the ratio of organic components to inorganic components in PM2.5.

### Outlook on Further Experimental Work

Additional sample analyses are currently ongoing and chemical composition data will be available for more time periods.

The chemical analysis will be extended to other organic molecules that are used as tracers for specific carbonaceous aerosol sources including biogenic (plant and bacterial materials) as well as anthropogenic (vehicle emissions) or mixed (biomass burning).

Additional data analysis will be performed by combining organic and inorganic/metal aerosol data (see companion poster, Aurelie Marcotte et. al.)