

Geographic Patterns of Mercury Deposition using the Lichen *Xanthoparmelia* in Maricopa County, Arizona, USA

K. Sweat, T. Zambo, and T. Nash III

School of Life Sciences, Arizona State University, PO Box 874501, Tempe AZ 85287-4501



A typical specimen of *Xanthoparmelia* sp. used for air pollution biomonitoring.

ABSTRACT

Geographical patterns of atmospheric metal deposition in Maricopa County, Arizona, were assessed using 2006 collections of the epilithic lichen *Xanthoparmelia* spp. This study is a re-sampling of the 28 locations used in an earlier investigation by Zschau and Nash in 1998, along with two new locations sampled to increase spatial resolution. Lichens were removed from the rock substrate, cleaned and homogenized. Initially the homogenized samples were analyzed for mercury [Hg] with a mercury analyzer and will be wet digested and analyzed by high resolution ICP-MS for a suite of up to 40 elements. Spatial Hg patterns are plotted with ArcGIS software to identify potential "hot spots". Potential sources of these high Hg levels, such as power generation facilities, were sought at or in the vicinity of these hot spots.

INTRODUCTION

Assessment of long-term air pollution patterns can often be difficult. Monitoring of pollutants over time can be resource intensive, and data collection with sampling devices needs to start years before analysis can begin. The use of living organisms as biomonitors is one method to overcome some of the drawbacks of monitoring air pollution over extended time periods. As long-lived, slow-growing organisms, lichens are useful as surrogate receptors in atmospheric deposition monitoring investigations, where the integration of long-term signals requires monitoring (Nash 1989, Garty 2001). Because they do not possess nutrient absorbing roots, as found in vascular plants, they have a major dependence on atmospheric sources of nutrients (Nieboer et al. 1978, Nieboer and Richardson 1980). Compared to soil nutrient pools, atmospheric concentrations of nutrients are quite low, and consequently nutrient concentrating mechanisms, such as particulate trapping (Garty et al. 1977), uptake to cell wall exchange sites or transport intracellularly (Beckett and Brown 1984; Brown and Beckett 1985), sequestering in complexes formed with lichen secondary metabolites (Purvis et al. 1987), or impaction of aerosols (Knops et al. 1996) are characteristic of lichens. As a consequence, lichens have often been used to document atmospheric deposition of radionuclides (e.g. Palmer et al. 1965, Seaward et al. 1988, Blazrov 1994) and various other atmospheric pollutants (Puckett 1988, Nash and Gries 1995). When an appropriate stratified sampling design (e.g. with respect to lichen species choice, microhabitat characteristics, atmospheric exposure, etc.) is employed, then both local and regional deposition patterns are readily discerned (Brutey 1993, Loppi and Bargagli 1996, Muir et al. 1993, Nash 1996), although care must be taken in the assessment of baseline levels (Gough et al. 1988; Bennett 2000). In an earlier investigation (Zschau et al. 2003), we determined past spatial patterns of atmospheric deposition across Maricopa County, Arizona, in 1998, based on samples from 28 sites, as analyzed by ICP-MS. The county is approximately L-shaped, extending over 200 km along its two longer axes, and contains the Phoenix metropolitan area, one of the fastest growing urban regions in the world. Although heavy industry is minimal within the Phoenix area, Arizona has historically been an important source of copper. Major sources of spatial elemental variation included copper mining and smelting (in an adjacent county), anthropogenic sources associated with the urban center (e.g. lead), and location of special geological features, such as mafic rocks with elevated concentrations of Co, Cr, Ni, and Sc relative to average abundances in the Earth's crust. In addition, temporal comparisons for six of the same sites within the county were possible, based on collections made in the mid-1970's (Nash et al. 2003), where sufficient lichen material was available for analysis. Decreases in copper (cessation of smelting) and lead (switch to unleaded gasoline) were demonstrable and increases in zinc were found. Other research has demonstrated that trace metals are detected in lichens near a coal power plant (Olmez et al., 1985) and in lichens transplanted to the region of a coal power plant (Garty & Hagemeyer, 1986).

METHODS AND GOALS



The overall objective is to document the spatial pattern of past elemental deposition as reflected in lichens (*Xanthoparmelia* spp.) as of 2006 within the region encompassing the greater metropolitan Phoenix area (Maricopa county) and, where possible, to determine historical trends in comparison to previous work. The genus *Xanthoparmelia* is selected as the most suitable biomonitor of metal deposition in metropolitan Phoenix region, because it is one of the few macrolichens readily obtaining enough material for analysis (Nash et al. 1977), is easily recognizable in the field, and has already been used for similar investigations (Zschau et al. 2003; Nash et al. 2003). Spatial patterns of atmospheric deposition of trace elements to these epilithic lichens will be assessed using the locations of the Zschau et al. (2003) study with two additional sites added to this research.

The lichen material will be cleaned and homogenized to prepare for metal analysis. Mercury content has been measured using a cold vapor mercury analyzer. The samples are to then be wet digested and analyzed by HF-ICP-MS for a suite of elemental concentrations (antimony [Sb], cadmium [Cd], cerium [Ce], chromium [Cr], cobalt [Co], copper [Cu], dysprosium [Dy], europium [Eu], gadolinium [Gd], gold [Au], holmium [Ho], lead [Pb], lutetium [Lu], neodymium [Nd], nickel [Ni], palladium [Pd], platinum [Pt], praseodymium [Pr], samarium [Sm], scandium [Sc], silver [Ag], terbium [Tb], thulium [Tm], tin [Sn], uranium [U], vanadium [V], ytterbium [Yb], yttrium [Y], and zinc [Zn]).

Surface maps for concentrations of at least mercury, cadmium, lead, copper, nickel, and zinc will be interpolated among the 30 locations using ArcGIS Geostatistics and Spatial Analyst packages. Multivariate statistical analysis will be used to analyze and correlate deposition patterns of the various metals. Because the *Xanthoparmelia* grows on rocks, part of the elemental variation observed in the area will doubtlessly be related to underlying variation in geology and associated blowing dust. Accordingly, it will be necessary to interpret the results in terms of basic knowledge of geochemistry (e.g. Levinson 1974; Taylor and McLennan 1985) as well as specific knowledge of the geochemistry in the region (e.g. Reynolds 1988; Titey and Anthony 1989). Because known pollution sources are present in the region, the results will also have to be interpreted in light of known emission data (e.g. U.S. Environmental Protection Agency 1997).

LITERATURE CITED

- Beckett, R.P., Brown, D.H., 1984. The relationship between cadmium uptake and heavy metal tolerance in the lichen genus *Peltigera*. *New Phytologist* 97, 301-311.
- Bennett, J.P., 2000. Statistical baseline values for chemical elements in the lichen *Hypogymnia physodes*. In: Agrawal, S.B., Agrawal, M. (Eds.), *Environmental Pollution and Plant Responses*. Lewis, Boca Raton, pp.343-353.
- Blazrov, L.G., 1994. The radionuclides in lichen thalli in Chernobyl and east Ureals areas after nuclear accidents. *Phyton (Austria)* 34, 85-94.
- Brown, D.H., Beckett, R.P., 1985. Minerals and lichens: localisation and effects. In: Vozzoni, C., Brown, D.H., Legas, M.E. (Eds.), *Surface Physiology of Lichens*. Universidad Complutense de Madrid, Madrid, pp. 127-149.
- Entwistle, John, 2004. The Determination of Mercury in Microwave Digests of Foodstuffs by ICP-MS. National Measurement System Valid Analytical Measurement (VAM) Program.
- Garty, J., 2001. Biomonitoring atmospheric heavy metals with lichens: theory and application. *Critical Review in Plant Sciences* 20, 309-371.
- Garty, J., 1985. Heavy Metals in the Lichen *Ramalina duriaei* Transplanted at Biomonitoring Stations in the Region of a Coal-Fired Power Plant in Israel After 3 Years of Operation. *Water, Air and Soil Pollution*, 38,311-323.
- Garty, J., Galun, M., Kessel, M., 1979. Localization of heavy metals and other elements accumulated in the lichen thallus. *New Phytologist* 82, 159-168.
- Knops, J.M.H., Nash, III, T.H., Schlessinger, W.H., 1996. The influence of epiphytic lichens on the nutrient cycling of an oak woodland. *Ecological Monographs* 66, 159-179.
- Levinson, A.A., 1974. *Introduction to Exploration Geochemistry*. Applied Publishing Ltd., Calgary, Canada.
- Lindsey, C.G., Chen, J., Dye, T.S., Richards, L.W., Blumenthal, D.L., 1999. Meteorological Processes Affecting the Transport of Emissions from the Navajo Generating Station to Grand Canyon National Park. *Journal of Applied Meteorology*, 38, 1031-1048.
- Loppi, S. and Bargagli, R., 1996. Lichen biomonitoring of trace elements in a geothermal area (central Italy). *Water, Air and Soil Pollution* 88, 1177-1187.
- McCune, B., Mefford, M.J., 1982. PC-ORD. Multivariate Analysis of Ecological Data, Version 4. MJM Software Design, Glenden Beach, Oregon, USA.
- Muir, D.E.G., Segstro, M.D., Wellbourn, P.M., Toom, D., Eisenreich, S.J., Macdonald, C.R., Weipold, D.M., 1993. Patterns of accumulation of airborne organochlorine contaminants in lichens from the Upper Great Lakes Region of Ontario. *Environmental Science and Technology* 27, 1211-1210.
- Nash, III, T.H., 1989. Metal tolerance in lichens. In: J. Shaw (ed.) *Metal Tolerance in Plants: Evolutionary Aspects*. CRC Press, Boca Raton, Florida, pp. 119-131.
- Nash, III, T.H., 1996. Nutrient, elemental accumulation and mineral cycling. In Nash III, T.H. (Ed.), *Lichen Biology*. Cambridge University Press, Cambridge, pp. 136-153.
- Nash, III, T.H., Gries, C., 1995. The use of lichens in atmospheric deposition studies with an emphasis on the Arctic. *Science for the Total Environment* 160/161, 729-738.
- Nash, III, T.H., Gries, C., Zschau, T., Getty, S., Ameron, Y., Zambrano, A., 2003. Historical patterns of metal atmospheric deposition to the epilithic lichen *Xanthoparmelia* in Maricopa County, Arizona, U.S.A. *Journal of Physics IV (France)* 107, 921-924.
- Nash, III, T.H., White, S.L., Marsh, J.E., 1977. Lichen and moss distribution and biomass in hot desert ecosystems. *Bryologist* 80, 470-479.
- Nieboer, E., Richardson, D.H.S., 1980. Lichens as monitors of atmospheric deposition. In: Eisenreich, S.J. (Ed.), *Atmospheric Pollutants in Natural Waters*. Ann Arbor Science, Ann Arbor, pp. 339-388.
- Nieboer, E., Richardson, D.H.S., Tomassin, F.O., 1976. Mineral uptake and release by lichens: an overview. *Bryologist* 81, 226-246.
- Olmez, I., Gulovaci, M.C., Gordoni, G.E., 1985. Trace Element Concentrations in Lichens Near a Coal-Fired Power Plant/Atmospheric Environment, 19, 1663-1669.
- Pacyna, E., Pacyna, J., 2002. Global Emission of Mercury from Anthropogenic Sources in 1995. *Water, Air and Soil Pollution* 127, 149-165.
- Palmer, H.E., Hanson, W.C., Griffin, B.J., Braby, L.A., 1965. Radioactivity measured in Alaskan natives, 1962-1964. *Science* 147, 620-621.
- Puckett, K.J., 1988. Bryophytes and lichens as monitors of metal deposition. *Bibliotheca Lichenologica* 30, 231-267.
- Purvis, O.W., Elix, J.A., Broomhead, J.A., Jones, S.C., 1987. The occurrence of copper-motile acid in lichens from copperiferous substrata. *Lichenologist* 19, 193-203.
- Ruick, J., Brill, T., Tates, D., 2002. Mercury in US coal: observations using the CD-AL/AL-ICR data. *Environmental Geology* 43, 247-259. online: <http://link.springer.com/link/service/journals/00254/bbs/3043003/30430271.htm>.
- Rand, A.E., 1998. Effects of a copper smelter on desert vegetation: a retrospective after 26 years. M.S. Thesis. Tempe: Arizona State University.
- Reynolds, S.J., 1988. *Geologic Map of Arizona*. Arizona Geological Survey Map 26, scale 1:1,000,000.
- Seaward, M.R.D., Heslop, J.A., Green, D., Byrnes, E.A., 1988. Recent levels of radionuclides in lichens from southwest Poland with particular reference to ¹³⁴Cs and ¹³⁷Cs. *Journal of Environmental Radioactivity* 7, 123-129.
- Shaw, J. E. 1995. Pattern recognition in lichens for source apportionment. *Atmospheric Environment* 29, 23-24.
- Taylor, S.R., McLennan, S.M., 1985. *The Continental Crust: Its Composition and Evolution*. Blackwell Scientific Publications, London.
- Titey, S.R., Anthony, E.Y., 1989. Laramide mineral deposits in Arizona. In: Jenny, J.P., Reynolds, S.J. (Eds.) *Geologic Evolution of Arizona*. Arizona Geological Society Digest 17, Tucson, pp. 485-514.
- U.S. Environmental Protection Agency, 2000. *Regional Air Quality Criteria for the Environmental Impact of Hazardous Air Polluting Sources*. EPA 440/9-00-001. U.S. Environmental Protection Agency, 1997. *Toxic Release Inventory 1987-1995*. Pollution Prevention and Toxics (PPT). EPA 749-C-97-003.
- Zschau, T., 1999. Spatial and temporal air pollution patterns of selected heavy metals in Maricopa County, Arizona. M.S. Thesis. Arizona State University, Tempe.
- Zschau, T., Getty, S., Gries, C., Ameron, Y., Zambrano, A., Nash III, T.H., 2003. Historical and current atmospheric deposition to the epilithic lichen *Xanthoparmelia* in Maricopa County, Arizona. *Environmental Pollution* 125: 21-30.



From Fig. 3. Spatial distribution of metal concentration as measured in *Xanthoparmelia* spp. in 1998. Zschau et al., 1999.

View of the Phoenix skyline from South Mountain Park.