



The stoichiometry of urban pests: nutrient composition in an arthropod predator-prey system across metropolitan Phoenix



Patricia Trubl¹ and J. Chadwick Johnson²

¹School of Life Sciences, Arizona State University and ²Division of Mathematical & Natural Sciences, Arizona State University at the West Campus.

INTRODUCTION

- Understanding the impacts of ‘human-induced rapid ecological change’ (HIREC) has recently been described as a ‘grand challenge’ for ecologists¹. Given that half the world’s population now resides in urban centers, urbanization is a prevalent example of HIREC that deserves study².
- Urbanization often results in the loss of native species diversity at the expense of a few urban-adapted taxa also known as ‘urban exploiters’³. However, it remains unclear why urban exploiters thrive following human disturbance, while others species are lost.
- The western black widow spider, *Latrodectus hesperus*, and its most common prey item in urban Phoenix, the decorated cricket (*Grylloblatta campodeiformis*) thrive in disturbed, urban habitat—often forming dense aggregations (i.e. infestations). Recently, we have shown that the average body mass and population density of urban *L. hesperus* are well predicted by their foraging success on *G. sigillatus*⁴. Furthermore, *L. hesperus* average body mass, population density and prey abundance vary significantly across urban Phoenix aggregations. As such, we were interested to learn more about the mechanisms underlying this predator-prey relationship among urban pests.
- Specifically, ecological stoichiometry (ES) allows for the quantification of multiple chemical elements and the balance of energy in ecological interactions⁵. Currently, in urban ecology, ES is used to understand how urbanization alters food web dynamics⁶.
- Here we test the hypothesis that urban habitats are spatially heterogeneous. Thus, we predict the carbon-nitrogen (C: N) stoichiometry of *L. hesperus* and *G. sigillatus* will show high levels of spatial variation. Moreover, we predict that *G. sigillatus* C: N stoichiometry shapes *L. hesperus* C:N stoichiometry across 10 urban sites within the greater metropolitan Phoenix, AZ area.

METHODS

- Urban collection sites (see Fig 1) were chosen if they met the following criteria: 1) sites had to be a minimum of 8 km apart and 2) sites had to contain a minimum of 10 female widows within 5000 m² (i.e. infestations). All sites were commercial subhabitats with xeric landscaping.
- L. hesperus* webs were located visually and all penultimate and adult females were captured from a site. *G. sigillatus* were collected within a 10m² area of any *L. hesperus* web located at a site. Crickets and spiders were collected from mid-August to mid-September 2010. Samples were stored at -20°C for later chemical analysis.
- All samples were dried at 60°C for 120 hours. *G. sigillatus* were ground into a homogenous powder using a ball mill (*Spex-Certiprep 8000D*) and analyzed for % C and % N using a *Perkins-Elmer 2400 CHN analyzer*. *L. hesperus* were cut longitudinally, and a subsample was analyzed for % C and % N using a *Perkins-Elmer 2400 CHN analyzer*.



Fig 1. Black widow aggregations studied across Phoenix, Arizona (Map from Trubl *et al.* 2012).

RESULTS

Nutrient composition proved to be variable between sites in terms of the C: N ratio ($F_{9,119} = 6.859, P < 0.001$, see Fig. 2a) and % N of black widows ($F_{9,119} = 5.157, P < 0.001$), and the C: N ratio ($F_{9,99} = 2.123, P = 0.035$, Fig. 2b) and the % N of the decorated cricket ($F_{9,99} = 2.07, P = 0.04$). High C: N ratio of the decorated cricket was significantly associated with increased % N of black widows across urban Phoenix ($F_{1,9} = 6.512, P = 0.034$, Fig. 3).




Fig 2. A. C: N ratio of black widows and B. the decorated cricket across 10 urban sites across metropolitan Phoenix, AZ.




Fig 3. *G. sigillatus* C:N ratio is a strong predictor of *L. hesperus* % N across urban Phoenix, AZ.

DISCUSSION

- We found strong site variation in C: N ratios similar to that described by Trubl *et al.* (2012) for widow population ecology parameters⁴. Because that previous study also found black widow foraging success on *G. sigillatus* was a significant predictor of spider mass and population density, we were surprised here to find very few correlations between cricket and spider nutrient composition.
- The one relationship documented showed black widow % N increase as cricket (prey) C: N increases. This rather perplexing result suggests to us that widows are deriving N from a different prey source.
- Alternatively, arthropod nutrient composition across Phoenix could be a product of genetic differences across urban sites. However, preliminary results in a population genetic study of black widows, including all sites within this study, indicate no genetic clustering by sites (i.e. high gene flow between these sampling locales).
- Future work will include the addition of Phosphorus (P) content of the black widows and decorated crickets across the sites studied herein. As P limitation increases with increasing trophic levels⁷, generalist arthropod predators, such as *L. hesperus* that experience high prey abundance but lower prey diversity (i.e. less nutrient diversity) characteristic of urban environments, may face strong P limitation.
- Multiple prey types will be included in future analyses to better represent the widows complete diet and better describe urban Phoenix’s arthropod food web.

LITERATURE CITED

- Sih A, *et al.* 2010. *Integrative and Comparative Biology* 50: 945-947
- Population Division of the Department of Economic and Social Affairs of the United Nations Secretariat, *World Population Prospects: The 2006 Revision and World Urbanization Prospects: The 2007 Revision*, <http://esa.un.org/unup>.
- Blair RB 1996 *Ecological Applications* 6: 506–519
- Trubl P, *et al.* 2012. *Urban Ecosystems* DOI 0.1007/s11252-011-0223-2
- Sterner, RW & Elser, JJ 2002. Princeton University Press.
- Singer GA & Battin TJ 2007. *Ecological Applications* 17: 376-389
- Elser JJ *et al.* 2000. *Nature* 408: 578-580

ACKNOWLEDGMENTS

We would like to thank Lindsay Miles and Theresa Gburek for aiding in arthropod collection. This material is based upon work supported by the National Science Foundation under grant no. DEB-0423704 CAP LTER.