

Conceptual Methods for Defining Mammalian Functional Traits in Urban Landscapes

Katherine C. B. Weiss^{1,2}, Courtenay A. Ray¹, Jan Schipper², and Sharon J. Hall¹ ¹School of Life Sciences, Arizona State University, ²Arizona Center for Nature Conservation, Phoenix Zoo





Background

Cities have been described as novel ecosystems that support unique assemblages of wildlife^{1,2}. Yet undefined are mechanisms for how species filter into cities to form communities². Functional traits that respond to people and the environment have been postulated as likely mechanisms for urban community assembly². However, significant disagreement exists on how to measure and compare functional traits across taxa³. Other approaches instead categorize species by their ability to avoid, utilize, or dwell within cities⁴. However, classifications fail to account for how species change across space and time⁵.

How do methods for the classification of species in cities differ, and how might we synthesize approaches to support the creation of more useful datasets?

We present a novel functional trait framework to be applied across terrestrial mammals within CAP LTER. We predict mammals closer to urban elements share physiological, behavioral, and cultural functional traits.

Challenges with Functional Traits

Many use functional traits to quantify species distributions in cities^{6,7,8}. Despite efforts to untangle how traits correspond with feedbacks between organisms and their environment:

- Functional trait studies often do not select traits based on the research question or community assembly processes
- Definitions remain unaligned (Table 1), limiting the comparability of current work across contexts³

Table 1. Example "functional trait" definitions in foundational functional ecology literature. Citations were determined on Google Scholar (https://scholar.google.com/) on November 5, 2018. Adapted from

Weiss and Ray (In Review) 3.				
	Publication	Defines functional traits as	Citations	
	Tilman (2001) ⁹	" organismal traits that influence one or more aspects of the functioning of an ecosystem"	547	
	Díaz and Cabido (2001) ¹⁰	" the characteristics of an organism relevant to its response to the environment and/or its effects on ecosystem functioning"	2169	
	Lavorel and Garnier (2002) ¹¹	" Response groups and effect groups Physiological, harder traits at the individual level are more commonly used for effect groups Whereas response groups are identified through community-level studies of changes in soft, morphological or behavioural traits in response to abiotic or biotic factors"	2025	
	McGill et al. (2006) ¹²	" a well-defined, measurable property of organisms, usually measured at the individual level and used comparatively across species that strongly influences organismal performance"	2451	
	Petchey and Gaston (2006) ¹³	" components of an organism's phenotype that influence ecosystem level processes"	1401	
	Violle et al. (2007) ¹⁴	"Any trait which impacts fitness indirectly via its effects on growth, reproduction and survival"	1881	

Limitations of Other Methods

An alternative to using functional traits is categorizing species as urban avoiders, utilizers, or dwellers⁵. However, categorizations have recently been called into question, as they fail to account for how species change over space, through time, and across heterogeneous environments⁴ (Fig. 1).




Figure 1. Illustrating the limitations of current categorizations of urban wildlife. Many classify species as urban avoiders, utilizers, or dwellers^{4,5}. However, **(A)**

categorizations may differ across cities depending upon regional species pools, and (B) the role a species plays within a city depends upon local landscape characteristics, and so may not be consistent⁵.

Unifying Approaches

functional traits for urban

left side is an adaptation of the

hierarchical filtering framework


determined by socio-ecological

interactions that filter through

urban species pools are

and Ray In Review)3.

We provide a unified definition of functional traits to support application across contexts and along urban gradients (Fig. 2). By focusing on aspects of an organism's physiology and behavior that correspond with fitness, we can ascertain longer-term trends in community assembly.



In applying our framework to terrestrial mammals in CAP LTER (Table 2), we advocate using traits that:

- Are resilient to genetic drift
- Have selective advantages for urban exploitation
- Support dispersal into the urban matrix
- May be selected for or against by human decision-making at the local scale

Mammal Traits for CAP LTER

Table 2. Mammalian functional traits of interest in CAP LTER. Presented traits are those expected to be important for community assembly processes and human preferences along urban gradients.

Trait	Corresponds with	Predicted Trends with Urbanization	
Activity pattern	Behavioral thermoregulation and human avoidance	Increased nocturnality ¹⁶	
Body size	Heat dissipation, human avoidance, and the increase of mesopredators in cities ¹⁷	Small to moderate body sizes	
Dentition and nutritional physiology	Urban resource use	Generalized dentition with few dietary restrictions	
Home range plasticity	Adaptability to changing patch sizes	Higher home range plasticity	
Behavioral plasticity	Acclimation to humans and increased species densities in urban areas ¹⁸	Greater behavioral plasticity	
Social structure	Population size and whether or not individuals collect in groups or individually	Solitary social behavior	
Facial structure and dentition	Human perception of wildlife as "cute" and non-threatening	Human preferred facial features (e.g., small and round)	

Conclusions & Future Directions

- As our work considers both adaptations and socio-ecological interactions, we believe our approach may support better predictions for how species assemble in heterogeneous landscapes
- We will use our framework in CAP LTER to identify how urbanization influences wildlife in Phoenix, AZ
- Our research will test a novel functional trait approach for the assessment and conservation of urban ecological communities across the U.S.



References

¹Grimm, N., et al. 2008. Science 319(5846): 756-760. ²Aronson, M., et al. 2016. Ecology 87(11): 2952-2963. ³Weiss, K.C.B., and C. A. Ray. In Review. Ecography. ⁴Fisher, J.D., et al. 2015. Conservation Biology 29(4): 1246-1248. ⁵Soulsbury, C., and P. White. 2015. Wildlife Research 42(7): 541-553. ⁶Devictor, V., et al. 2007. Conservation Biology 21:741–751. ⁷Vandewalle, M., et al. 2010. Biodiversity and Conservation 19:2921–2947. ⁸Santini, L., et al. 2018. Ecology Letters:1–12. ⁹Tilman, D. 2001. Encyclopedia of Biodiversity, pgs. 109-120. ¹⁰Díaz, S., and M. Cabido. 2001. Trend in Ecology & Evolution 16(11): 646-655. ¹¹Lavorel, S., and E. Garnier. 2002. Functional Ecology 16:545–556. ¹²McGill, B.J., et al. 2006. Trends in Ecology & Evolution 21(4): 178-185. ¹³Petchey, O.L. and Gaston, K.J. 2006. Ecology Letters. 9: 741–758. ¹⁴Violle, C., et al. 2007. Oikos 116(5): 882-892. ¹⁵Vellend, M. 2016. The theory of ecological communities. Princeton University. ¹⁶Gaynor, K.M. et al. 2018. Science. 360: 1232–1235.

¹⁷Crooks, K.R., and M.E. Soulé. 1999. Nature 400:563–566

¹⁹Borgi, M., et al. 2014. Frontiers in Psychology. 5: Article 411.

¹⁸Shochat, E., et al. 2006. Trends in Ecology & Evolution 21:186–191.